



Castion®



seasonality

Early-Mid primocane variety,
15 days later than Enrosadira®.

Early-Mid florican variety,
10 days later than Enrosadira®.



climatic areas

It performs better in **areas with at least 400 chill hours**, but it is also adapted to low chill areas where the primocane-fruited season can be extended.



plant structure and agronomic description

Long and erect canes, medium amount of spines, dense foliage. Variety suitable for long canes production. Abundant bloom on lateral shoots of medium size, good flower pollination. Auto-fertile variety, but the presence of pollinator insects is always helpful.



yield

Productive plant, specially after 1 year of transplant.
Easy to pick.



berry description

Firm, large, conical and **shiny** red berries, with good shelf life. Great flavor: **sweet- acid** and **aromatic**.



RIPENING PERIOD																												
ENROSADIRA®																												
HALLEY																												
CASTION®																												
DOROTEA																												
AURORA®																												
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
	May				June				July				August				September				October							

Data collected from Vivai Molari in Trento (2018/2019)

VARIETY	SEASONALITY	CHILL REQUIREMENT	BERRY WEIGHT (g)	BERRY COLOR*
ENROSADIRA®	Early	Mid - High	6	light red
HALLEY	Early	Mid - High	5,3	red
CASTION®	Early - Mid	Mid - High	6,6	shiny red
DOROTEA	Mid - Late	Mid - High	7,1	red
AURORA®	Late	Mid - High	5,7	pink red
POLKA	Early	High	6,6	red
AUTUMN BLISS	Early	High	4	red

*BERRY COLOR at harvest day